

Polyphenols and prevention of cardiovascular diseases

Claudine Manach, Andrzej Mazur and Augustin Scalbert

Purpose of review

Polyphenols are the most abundant dietary antioxidants and research on their role in the prevention of degenerative diseases has developed quickly over these last few years. This paper reviews the recent studies on the prevention of cardiovascular diseases by polyphenols, focusing on human studies.

Recent findings

A large number of recent intervention studies have shown that several biomarkers of cardiovascular risk are influenced by the consumption of polyphenol-rich foods. Effects on biomarkers of oxidative stress, lipemia and inflammation appear so far inconclusive. More-consistent effects have been observed on endothelial function and haemostasis and support a reduction of risk by polyphenols in agreement with the few epidemiological studies already published. All clinical studies have used foods or beverages containing a mixture of different polyphenols and the exact nature of the most active compounds remains largely unknown. Absorption, metabolism and elimination vary widely between polyphenols. These data on bioavailability should be taken into account to improve the experimental design and the interpretation of the observed effects.

Summary

Future intervention studies should include a detailed assessment of the bioavailability of polyphenols. Beyond clinical trials carried out with polyphenol-rich foods, more studies with pure polyphenols will also be needed to establish their role in the prevention of cardiovascular diseases.

Keywords

antioxidant, bioavailability, cardiovascular diseases, clinical trials, endothelial function, flavonoids, haemostasis, lipemia, polyphenols

Introduction

Polyphenols are common constituents of the human diet, present in most foods and beverages of plant origin. They are considered to contribute to the prevention of various degenerative diseases, including cardiovascular diseases. This assumption originally came from *in vitro* studies, showing the antioxidant properties of several polyphenols and their ability to modulate the activity of various enzymes. More evidence for a protective role of polyphenols against cardiovascular diseases arose from a number of epidemiological studies, clinical trials, experiments on animal models and mechanistic studies. However, data are still often contradictory. Two reasons in particular can be inferred. The first one is that the polyphenol family encompasses very diverse compounds with highly different bioavailabilities. Hence the results obtained for one polyphenol cannot be generalized to others. The second point is that polyphenols are now known to be largely metabolized in the body and native compounds most often tested in *in vitro* studies are virtually absent in the tissues. The relevance of these *in vitro* studies with respect to the reduction of cardiovascular risk must therefore be questioned.

This paper reviews the recent studies on the prevention of cardiovascular diseases by polyphenols, focusing on *in vivo* studies. Results are discussed in the light of our knowledge on bioavailability.

Polyphenols in animal models of atherosclerosis

The effect of polyphenol-rich foods or polyphenol extracts on the development of atherosclerotic lesions has been studied in apolipoprotein E^{-/-} mice and hamsters. A reduction of the lesion was observed upon consumption of a tea extract [1], green and black tea [2], pomegranate juice [3], grape extracts [4], red wine, dealcoholized wine and wine polyphenols [4,5,6*]. The atheroprotective effect of dietary soy isoflavones in apolipoprotein E^{-/-} mice was shown to require the presence of estrogen receptor- α and was unrelated to plasma cholesterol [7**]. Only a few studies have evaluated and demonstrated an antiatherosclerotic effect of pure polyphenols (for example quercetin and catechin) [8]. The reduction of atherosclerotic lesion by polyphenol intake is not necessarily related to the modifications in the oxidative stress biomarkers and lipid parameters. This suggests that other mechanisms, for example the anti-inflammatory action of polyphenols, could be involved. These animal experiments suggest a protective

Curr Opin Lipidol 16:000–000. © 2005 Lippincott Williams & Wilkins.

Unité des Maladies Métaboliques et Micronutriments, INRA, 63122 Saint-Genès Champanelle, France

Correspondence to Augustin Scalbert, Unité des Maladies Métaboliques et Micronutriments, INRA, 63122 Saint-Genès Champanelle, France
Tel: +33 473 62 47 87; fax: +33 473 62 46 38; e-mail: scalbert@clermont.inra.fr

Current Opinion in Lipidology 2005, 16:000–000

Abbreviations

FMD flow-mediated dilation
HDL high-density lipoprotein
LDL low-density lipoprotein

© 2005 Lippincott Williams & Wilkins
0957-9672

effect of polyphenols but they cannot be extrapolated directly to humans. In particular, the doses applied are often higher than those to which humans are commonly exposed with their diet.

Polyphenols and cardiovascular diseases in epidemiological studies

Various epidemiological studies have shown an inverse association between the consumption of polyphenols or polyphenol-rich foods and the risk of cardiovascular diseases. A meta-analysis including seven case-control and 10 cohort studies suggested a reduction of the risk of myocardial infarction of 11% upon consumption of three cups of tea per day [9]. One more recent prospective study and three cross-sectional studies tend to confirm these protective effects of tea against cardiovascular risk [10–13].

A moderate wine consumption has consistently been associated with a reduced risk of cardiovascular diseases. However, the respective contribution of alcohol and polyphenols to these effects was not well clarified. A meta-analysis based on 26 cohort and case-control studies showed that light to moderate intake of both wine and beer was associated with a reduction of vascular risk [14[•]]. The reduction was more pronounced for wine, possibly due to its higher polyphenol content. Two recent cohort studies in which wine was compared to beer and spirits or red wine compared to white wine also suggested that wine, and more particularly red wine, which is about 10 times richer in polyphenols than white wine, were more protective than beer or spirits [15[•]]. An inverse association between ischemic stroke and wine but not beer and spirit consumption was also observed in a prospective study [16].

Correlation between the intake of flavonols, flavones, catechins and lignans and the risk of coronary heart diseases and stroke have been looked for [17]. Out of 18 prospective studies on coronary heart diseases or stroke, 10 showed an inverse association between intake and risk, and the other no association. A cross-sectional study showed a reduction of cardiovascular risk in US postmenopausal women consuming the largest amounts of isoflavones [18]. More studies are needed, in particular on the more abundant anthocyanins, proanthocyanidins and phenolic acids polyphenol classes. They have not been studied so far due to the lack of appropriate food-composition tables or their too-recent publication.

Clinical studies

Altogether, animal experiments and epidemiological studies suggest a protective role of polyphenols. Several mechanisms of action have been proposed and explored in clinical studies (Table 1). Most intervention studies in humans have been carried out with polyphenol-rich foods

Table 1. Proposed mechanisms by which polyphenols may reduce risk for cardiovascular diseases

Antioxidants	Scavenge reactive oxygen and nitrogen species
	Chelate redox-active transition metal ions
	Spare and interact with other antioxidants
	Inhibition of the redox-sensitive transcription factors
	Inhibition of pro-oxidant enzymes
	Induction of antioxidant enzymes
Growth of atherosclerotic plaque	Reduce adhesion molecule expression
	Anti-inflammatory
	Reduce the capacity of macrophages to oxidatively modify LDL
Platelet function and haemostasis	Inhibit platelet aggregation
Blood pressure and vascular reactivity	Promote nitric oxide-induced endothelial relaxation
Plasma lipids and lipoproteins	Reduce plasma cholesterol and triglycerides

LDL, low-density lipoprotein.

and in particular tea, wine, cocoa or soy. These foods and beverages do not only contain polyphenols and only the properly controlled studies are considered here (soy with or without isoflavones, tea versus caffeine, wine versus alcohol).

Antioxidant effects of polyphenols

Antioxidant effects of polyphenols have been reviewed elsewhere [19,20]. The most recent human intervention trials are discussed here (Table 2). When the source of polyphenols was consumed for 1–12 weeks, an increase in the plasma antioxidant capacity or in the concentration of antioxidants such as vitamin E, vitamin C, β -carotene and uric acid was observed in some studies, whereas no change were observed in other studies. The same contrasting results were obtained for lipid-oxidation products in plasma and low-density lipoprotein (LDL) oxidizability measured *ex vivo*. These discrepancies could be explained by kinetic effects, polyphenols being largely eliminated in the 2 to 8 hours following ingestion [21], and therefore virtually absent in fasting plasma samples collected after a night washout.

In acute studies, polyphenols are ingested alone or with a meal and blood samples are collected in the few hours following ingestion. Coffee, red wine and grape-seed extract all increased the postprandial plasma antioxidant capacity (Table 2). Olive oil showed no effect, probably due to the phenol intake being too low. A decrease of the level of plasmatic F₂-isoprostane, a marker of lipid oxidation, was also observed after ingestion of a cocoa drink [22[•]]. This last result is important as it shows a diminution of lipid damage shortly after the consumption of polyphenols. In these acute studies, the maximum plasma antioxidant capacity is usually reached 1–4 h after ingestion of the polyphenol source. This delay is similar to the time needed to reach the maximal concentration in plasma for most antioxidant polyphenols [23[•]].

Table 2. Intervention studies with polyphenols and effects oxidative stress biomarkers

Food	Food quantity (polyphenol intake)	Control	Subjects (number)	Duration of intervention	Biomarkers affected	Biomarkers not significantly affected	Reference
Chronic studies							
Black tea+fried onions	300 ml/day+150 g/day (131 mg flavonoids/day)	Iso-caloric low-flavonoid diet	Healthy subjects (32)	1 week		Plasma F ₂ -isoprostane, LDL-MDA, vitamin C, vitamin E, β -carotene, iron	[27]
Green or black tea	1000 ml/day	Caffeine solution	Healthy subjects (13)	1–4 weeks		Urinary F ₂ -isoprostane	[28]
Black tea	5 servings/day	Caffeinated drink	Hypercholesterolemic patients (15)	3 weeks		Plasma AOC, oxidized LDL, LDL-TBARS, F ₂ -isoprostanes; urinary 8-OHdG	[29]
Red wine	250 ml/day	White wine, champagne	Healthy subjects (18)	3 weeks	Plasma AOC \uparrow	Plasma TBARS, uric acid, vitamin E	[30]
Dealcoholized red wine	375 ml	White wine, red wine	Smokers (18)	2 weeks	Plasma F ₂ -isoprostane \downarrow , urinary F ₂ -isoprostane \downarrow , serum γ -GT \downarrow , serum uric acid \downarrow , β -carotene \uparrow	Plasma vitamin C, vitamin E	[31]
Extra-virgin olive oil	20 g/day	Regular olive oil	Hyperlipidemic patients (10)	6 weeks	LDL oxidizability \downarrow	LDL-vitamin E	[32]
Virgin olive oil	25 ml/day	Refined olive oil	Healthy subjects (30)	3 weeks	Oxidized LDL \downarrow , LDL oxidizability \uparrow		[33]
High-phenol olive oil	70 g/day (18 mg phenols/day)	Low-phenol olive oil	Smokers (25)	3 weeks		Plasma AOC, MDA, lipid hydroperoxides, protein carbonyl	[34]
Cocoa powder+dark chocolate	22 g/day+16 g/day (466 mg proanthocyanidins/day)	No cocoa, no chocolate	Healthy subjects (23)	4 weeks	Serum AOC \uparrow , LDL oxidizability \downarrow		[35]
Cocoa extract	(234 mg flavanols/day)	Placebo	Healthy subjects (32)	4 weeks	Plasma vitamin C \uparrow , uric acid \uparrow	Plasma AOC, MDA, vitamin C, vitamin E, F ₂ -isoprostanes	[36]
Bilberries, lingonberries+ black currants	100 g/day	No berries	Healthy subjects (20), 60 years old	8 weeks	Serum vitamin C \uparrow	Serum AOC, vitamin E; LDL oxidizability	[37]
Acute studies							
Red wine	250 ml	No wine	Healthy subjects (15)	<1 day	Plasma AOC \uparrow , uric acid \uparrow		[38]
Red wine	300 ml	No wine	Healthy subjects (3)	<1 day	Plasma AOC \uparrow		[30]
Red wine	400 ml	Hydroalc. sol.	Healthy subjects (6)	<1 day	Plasma AOC \uparrow , vitamin E \uparrow , protein thiols \uparrow	Plasma vitamin C, uric acid	[39]
Grape-seed extract	300 mg	No grape seed extract	Healthy subjects (8)	<1 day	Plasma AOC \uparrow , lipid hydroperoxyde \downarrow	LDL oxidizability	[40]
Virgin olive oil	50 ml	Baseline	Healthy subjects (16)	<1 day	LDL oxidizability \downarrow	Plasma vitamin E	[41]
Fortified olive oil	47 g (31 mg phenols)	Olive oil with no phenols	Healthy subjects (12)	<1 day		LDL oxidizability	[42]
Cocoa drink	100 ml (187 mg flavanols)	Low-flavonoid cocoa drink	Healthy subjects (20)	<1 day	Plasma F ₂ -isoprostane \downarrow	Plasma AOC, MDA, vitamin C, vitamin E	[22*]

In the acute studies in the second half of the table, the source of polyphenols is ingested together with bread, breakfast or a meal. AOC, antioxidant capacity; γ -GT, γ -glutamyl transpeptidase; 8-OHdG, 8-hydroxy-2'-deoxyguanosine; LDL, low-density lipoprotein; MDA, malondialdehyde; TBARS, thiobarbituric acid-reactive substances.

Polyphenols thus affect transiently some markers of oxidative stress. However, the link between these effects and the risk of cardiovascular diseases remains to be established. A direct scavenging of free radicals by polyphenols, as often suggested, may not be the key mechanism explaining their effects on oxidative stress biomarkers and cardiovascular risk factors. Polyphenol concentrations in plasma are low (usually 1 μM or less) and much lower than those of uric acid, an endogenous antioxidant (150–450 μM). The extensive conjugation of polyphenols to glucuronic acid and sulfate groups in the tissues may markedly decrease their antioxidant capacity, as described for quercetin and isoflavones [24,25^{*}]. Red-wine polyphenols were shown to limit the development of atherosclerotic lesions in apolipoprotein E-deficient mice, independently of the lipid peroxidation [6^{*},26^{*}]. A better knowledge of the mechanisms involved in cell-mediated modulation of the redox status by polyphenol exposure is needed to better evaluate the significance of these antioxidant effects in the prevention of cardiovascular diseases.

Polyphenols and lipemia

Out of 12 recent intervention studies with different polyphenol sources lasting 1–13 weeks, six showed no effects on lipids in plasma (Table 3). In the other six studies, polyphenol ingestion resulted in a decrease of the level of total cholesterol, LDL-Cholesterol, apolipoprotein B or lipoprotein (a), or an increase of the level of high-density lipoprotein (HDL)-cholesterol and apolipoprotein A-I. Such an improvement in the lipid parameters were observed after ingestion of polyphenols from tea, virgin olive oil, cocoa, soybean or red clover. Both normo- and hypercholesterolemic subjects showed improvement or no change in the lipid parameters.

Six trials were carried out with either soy proteins differing in their isoflavone contents or with isoflavone extracted from red clover against placebo (Table 3). Three of these studies showed an improvement of some lipid parameters. In a recent meta-analysis of 10 human trials, soy proteins were shown to effectively reduce LDL-Cholesterol and increase HDL-Cholesterol but the extent of these effects could not be related to the dose of isoflavones ingested together with the proteins [43^{**}]. In ovariectomized monkeys, the consumption of soy but not of isolated isoflavone reduced cholesterolemia [44]. More studies with isolated isoflavones are needed to establish the hypolipidemic effects of isoflavones.

Polyphenols and endothelial function

Dysfunction of the vascular endothelial cells can be measured by flow-mediated dilation (FMD) of the brachial artery. The large majority of studies testing red wine [51–53], grape juice [54], black tea [55,56], soy [57,58] or cocoa [59] have shown a net beneficial effect on FMD.

Table 3. Intervention studies with polyphenols and effects on plasma lipids

Food	Amount of food ingested (polyphenol intake)	Control	Subjects (number)	Duration of intervention	Lipid parameters affected	Lipid parameters not affected	Reference
Black tea + fried onions	300 ml/day + 150 g/day (131 mg flavonoids/day)	Isocaloric low-flavonoid diet	Healthy subjects (32)	1 week		Total fat, SFAs, PUFAs, MUFAs, C	[27]
Black tea	5 servings/day	Caffeinated drink	Hypercholesterolemic subjects (15)	3 weeks	C ↓, LDL-C ↓, apoB ↓, lipoprotein (a) ↓		[29]
Red wine	250 ml/day	White wine, champagne	Healthy subjects (18)	3 weeks		TG, C, PL, apoA-I, apoB	[30]
Dealcoholized red wine	375 ml	White wine, red wine	Smokers (18)	2 weeks		TG, C, LDL-C, HDL-C	[31]
Virgin olive oil	25 ml/day	Refined olive oil	Healthy subjects (30)	3 weeks	HDL-C ↑	TG, C, LDL-C	[33]
Cocoa powder + dark chocolate	22 g/day + 16 g/day (466 mg proanthocyanidins/day)	No cocoa, no chocolate	Healthy subjects (23)	4 weeks	HDL-C ↑	LDL/HDL	[35]
Isoflavone-rich soy protein	63 g/day (123 mg isoflavones/day)	Isoflavone-poor soy protein	Hypercholesterolemic (36)	6 weeks		LDL size, C concentration in LDL	[45]
Isoflavone-rich soy protein	25 g/day (50 mg isoflavones/day)	Isoflavone-poor soy protein	Hypercholesterolemic (42)	6 weeks		TG, C, LDL-C, HDL-C, apoB, apoA-I	[46]
Isoflavone-rich soy protein	(56 mg isoflavones/day)	Isoflavone-poor soy protein	Healthy subjects (22)	17 days	HDL-C ↑, apoA-I ↑	TG, C, LDL-C, apoA-II, apoB	[47]
Isoflavone-rich soy protein	(132 mg isoflavones/day)	Isoflavone-poor soy protein	Healthy women (18)	13 weeks	LDL-C ↓	TG, C, HDL-C, apoA-I, apoB, lipoprotein (a)	[48]
Red clover extract	(86 mg isoflavones/day)	Placebo	Premenopausal women (12)	12 weeks		TG, C, LDL-C, HDL-C, lipoprotein (a)	[49]
Biochanin-rich red clover extract	(40 mg isoflavones/day)	Placebo	Healthy subjects (40)	6 weeks	LDL-C ↓	TG, C, HDL-C	[50]

apo, apolipoprotein; C, cholesterol; HDL, high-density lipoprotein; LDL, low-density lipoprotein; MUFA, mono-unsaturated fatty acids; PUFA, polyunsaturated fatty acids; SFA, saturated fatty acids; TG, triglycerides.

These effects were observed in both healthy adults and coronary patients.

The positive effect of wine should be attributed to polyphenols rather than to alcohol. Dealcoholized red wine showed similar effect compared to whole red wine, whereas vodka did not affect the FMD of healthy volunteers [51,52]. Purple grape juice was also reported to improve the FMD in patients with coronary heart disease [54]. On the other hand, Whelan *et al.* [53] showed that in spite of a 6-fold lower polyphenol content, white wine equally improved the impaired endothelium function observed in adults with coronary heart disease, compared to red wine. If this result is confirmed, it implies that the active molecules might not be proanthocyanidins or anthocyanins, which are characteristic of red wine.

Consumption of black tea has also been shown to improve FMD in 50 patients with coronary heart disease [56]. After acute or long-term (30-day) tea intake, FMD increased to reach values comparable to that of healthy volunteers. Improvement of FMD upon long-term consumption of black tea was also observed in subjects with mild hyperlipemia [60]. A preliminary study on 10 young healthy volunteers suggested that black tea may also improve coronary endothelial function, as measured by a non-invasive transthoracic Doppler echocardiography method [61].

Cuevas *et al.* [58] observed a significant improvement of FMD in postmenopausal women having mild hypercholesterolemia with a short (4-week) supplementation with soy isolate protein providing isoflavones at 80 mg/day. Two recent double-blind studies merit particular attention because they used pure genistein [57,62]. Consumption of a genistein supplement (54 mg/day) for 6 or 12 months improved FMD in healthy postmenopausal women compared to placebo, increased levels of nitrites/nitrates and decreased plasma endothelin-1 levels. Steinberg *et al.* [63] compared the effect of isolated soy proteins with high or low isoflavone contents in healthy postmenopausal women. After 6 weeks, a non-significant trend for increased vasodilatory response was observed in the high isoflavone group compared to the low-isoflavone and control groups.

Polyphenols and inflammation

Polyphenols have a variety of anti-inflammatory and immune-modulating effects that may be of relevance to atherosclerosis. Even if *in vitro* studies have extensively provided arguments for anti-inflammatory effects of various polyphenols, human studies provide only a few tangible results. Supplementation with black tea, green tea, green tea polyphenol isolate, red wine, grape juice and cocoa affected only a few blood parameters related to inflammation, such as circulating adhesion molecules,

cytokines and mediators [35,64^{*},65–67]. Soy isoflavone supplementation for 6 weeks did not significantly affect concentrations of circulating adhesion molecules in postmenopausal women [63]. However, reduction of circulating vascular cell-adhesion molecule (VCAM)-1 concentrations was observed after 6-week administration of formononetin (daidzein precursor)-enriched isoflavones [68]. The limited anti-inflammatory effects of polyphenols observed in these human studies undoubtedly result from the fact that they were carried out on healthy subjects and rarely studied immune cell functions.

Polyphenols and haemostasis

Supplementation of the diet of experimental animals with polyphenols is known to increase bleeding time and to reduce *ex vivo* platelet aggregation [20]. Similarly, different clinical trials carried out on healthy subjects have shown an improvement of haemostasis after acute or chronic (2–4 weeks) consumption of red wine, tea or cocoa powder. With one exception [69], all recent studies run with red wine, black tea or cocoa against proper controls showed an inhibition of *ex vivo* collagen- or ADP-induced platelet aggregation [36,70–72].

Bioavailability and bioefficacy of polyphenols

Bioavailability differs greatly between polyphenols. Some polyphenols like soy isoflavones are well absorbed through the gut barrier whereas others such as proanthocyanidins, abundant in wine or cocoa, or thearubignins, the main polyphenols in black tea, are hardly absorbed [21]. Furthermore, each polyphenol may be present in foods in different forms, which can affect the intestinal absorption. Quercetin glucosides from onions are much more efficiently absorbed than other quercetin glycosides such as rutin present in tea or apple [73]. The total polyphenol content of tested food sources, generally stated in clinical trials, should be thus completed by a precise description of their polyphenol composition. This may avoid misleading interpretations. For example, effects of proanthocyanidin-rich extracts are systematically attributed to the major polymers whereas minor compounds may be better absorbed and more efficient.

Determination of plasma concentrations or urinary excretion of the polyphenol metabolites may facilitate the interpretation of clinical studies. High inter-individual variability has been frequently reported in the efficiency of intestinal absorption and metabolism, especially where the gut microflora are involved. For example, Meyer *et al.* [74^{*}] recently reported that only the volunteers who were able to convert daidzein into the microbial metabolite equol (35% of the subjects), gained benefit from consumption of soy products with significant reductions of cholesterol, LDL-Cholesterol, plasma triglycerides and lipoprotein (a). No significant effect was observed when the whole group of subjects was considered.

The peak of absorption and the elimination rate have also been shown to differ markedly among polyphenols. Catechins which are rapidly absorbed and eliminated from plasma (within 4 h) should have a more transient impact than other compounds such as isoflavones, which have a higher life span in the arteries. This kinetic data, now known for most polyphenols, could usefully be correlated with the observed effects.

The doses applied in animal experiments or in *in vitro* studies often exceed those to which human tissues may be exposed. According to the maximum concentrations obtained in human plasma after nutritional intakes [21], all the effects observed with doses above 2 μ M are unlikely to occur *in vivo*. Furthermore, polyphenols are present in plasma as glucuronide and sulfate ester derivatives. Biological activities of the conjugated metabolites are known to be influenced by conjugation. For example the affinities of daidzein and genistein for estrogen receptors are 10 and 40 times higher than those of their respective glucuronides [75]. It is thus crucial that all studies on the mechanisms of action of polyphenols, using very sophisticated methods but non-physiological compounds, can be complemented by studies using the metabolites actually found in the body. Non-absorbed polyphenols such as proanthocyanidins should never be tested in cultured cells or isolated organs since they have no chances of reaching the inner tissue level as such. Unfortunately, very few studies used relevant metabolites and doses so that our knowledge of the mechanisms of action of polyphenols are still very limited [76**].

Conclusion

Several biomarkers of cardiovascular risk are influenced by the consumption of polyphenol-rich foods. Effects on biomarkers of oxidative stress, lipemia and inflammation appear inconclusive. More-consistent effects have been observed on endothelial function and haemostasis and support a reduction of risk by polyphenols in agreement with the few epidemiological studies so far published.

All clinical studies have used foods or beverages containing a mixture of different polyphenols and the exact nature of the most active polyphenols remains largely unknown. Some polyphenols are highly bioavailable whereas others are hardly absorbed through the gut barrier. Until we know more about the activity of pure polyphenols in humans and targets of action of circulating polyphenols, it remains difficult to predict their protective efficacy against cardiovascular diseases. The lack of effects of polyphenols in some studies could be explained by rapid elimination of polyphenols once absorbed. Indeed, their effects are often more easily seen in the postprandial stages rather than in fasting conditions as often done. Together with the nature of the most active polyphenols, optimal doses to prevent cardiovascular

diseases need to be determined to formulate reliable nutritional recommendations.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

- 1 Miura Y, Chiba T, Tomita I, *et al*. Tea catechins prevent the development of atherosclerosis in apolipoprotein E-deficient mice. *J Nutr* 2001; 131:27–32.
- 2 Vinson JA, Teufel K, Wu N. Green and black teas inhibit atherosclerosis by lipid, antioxidant, and fibrinolytic mechanisms. *J Agric Food Chem* 2004; 52:3661–3665.
- 3 Kaplan M, Hayek T, Raz A, *et al*. Pomegranate juice supplementation to atherosclerotic mice reduces macrophage lipid peroxidation, cellular cholesterol accumulation and development of atherosclerosis. *J Nutr* 2001; 131:2082–2089.
- 4 Auger C, Gerain P, Laurent-Bichon F, *et al*. Phenolics from commercialized grape extracts prevent early atherosclerotic lesions in hamsters by mechanisms other than antioxidant effect. *J Agric Food Chem* 2004; 52:5297–5302.
- 5 Vinson JA, Teufel K, Wu N. Red wine, dealcoholized red wine, and especially grape juice, inhibit atherosclerosis in a hamster model. *Atherosclerosis* 2001; 156:67–72.
- 6 Waddington E, Puddey IB, Croft KD. Red wine polyphenolic compounds
 - inhibit atherosclerosis in apolipoprotein E-deficient mice independently of effects on lipid peroxidation. *Am J Clin Nutr* 2004; 79:54–61.
 These results suggest that lipid deposition is independent of lipid oxidation and that the protective action of red-wine polyphenols is independent of any antioxidant action of these compounds.
- 7 Adams MR, Golden DL, Register TC, *et al*. The atheroprotective effect of
 - dietary soy isoflavones in apolipoprotein E^{-/-} mice requires the presence of estrogen receptor- α . *Arterioscler Thromb Vasc Biol* 2002; 22:1859–1864.
 The results indicate a necessary role for estrogen receptor- α -dependent processes in mediating the atheroprotective effects of dietary soy isoflavones.
- 8 Hayek T, Fuhrman B, Vaya J, *et al*. Reduced progression of atherosclerosis in apolipoprotein E-deficient mice following consumption of red wine, or its polyphenols quercetin or catechin, is associated with reduced susceptibility of LDL to oxidation and aggregation. *Arterioscler Thromb Vasc Biol* 1997; 17:2744–2752.
- 9 Peters U, Poole C, Arab L. Does tea affect cardiovascular disease? A meta-analysis. *Am J Epidemiol* 2001; 154:495–503.
- 10 Mukamal KJ, Maclure M, Muller JE, *et al*. Tea consumption and mortality after acute myocardial infarction. *Circulation* 2002; 105:2476–2481.
- 11 Mennen LI, Sapiño D, De Bree A, *et al*. Consumption of foods rich in flavonoids is related to a decreased cardiovascular risk in apparently healthy French women. *J Nutr* 2004; 134:923–926.
- 12 Hakim IA, Alsaif MA, Alduwaihi M, *et al*. Tea consumption and the prevalence of coronary heart disease in Saudi adults: results from a Saudi national study. *Prev Med* 2003; 36:64–70.
- 13 Hirano R, Momiyama Y, Takahashi R, *et al*. Comparison of green tea intake in Japanese patients with and without angiographic coronary artery disease. *Am J Cardiol* 2002; 90:1150–1153.
- 14 Di Castelnuovo A, Rotondo S, Iacoviello L, *et al*. Meta-analysis of wine and
 - beer consumption in relation to vascular risk. *Circulation* 2002; 105:2836–2844.
 This is the first meta-analysis suggesting a different impact of two alcoholic beverages on vascular risk.
- 15 Klatsky AL, Friedman GD, Armstrong MA, Kipp H. Wine, liquor, beer, and
 - mortality. *Am J Epidemiol* 2003; 158:585–595.
 One of the main interests of this large cohort study is to present results on different causes of mortality for three alcoholic beverages at different levels of consumption. This is one of the few epidemiological studies to address the differences between red and white wine.
- 16 Djousse L, Ellison RC, Beiser A, *et al*. Alcohol consumption and risk of ischemic stroke: the Framingham Study. *Stroke* 2002; 33:907–912.
- 17 Arts ICW, Hollman PCH. Polyphenols and disease risk in epidemiological studies. *Am J Clin Nutr* 2005 (in press).
- 18 de Kleijn MJ, van der Schouw YT, Wilson PW, *et al*. Dietary intake of phytoestrogens is associated with a favorable metabolic cardiovascular risk profile in postmenopausal U.S. women: the Framingham study. *J Nutr* 2002; 132:276–282.

- 19 Williamson G, Manach C. Bioavailability and bioefficacy of polyphenols in humans: II A review of 93 intervention studies. *Am. J. Clin. Nutr.* 2005 (in press).
- 20 Scalbert A, Manach C, Morand C, *et al.* Dietary polyphenols and the prevention of diseases. *Crit Rev Food Sci Nutr* 2005 (in press).
- 21 Manach C, Williamson G, Morand C, *et al.* Bioavailability and bioefficacy of polyphenols in humans: I- A review of 97 bioavailability studies. *Am J Clin Nutr* 2005 (in press).
- 22 Wiswedel I, Hirsch D, Kropf S, *et al.* Flavanol-rich cocoa drink lowers plasma F(2)-isoprostane concentrations in humans. *Free Radic Biol Med* 2004; 37:411–421.
This is one of the few acute studies demonstrating an effect on the oxidation of lipids rather than on their oxidizability.
- 23 Manach C, Scalbert A, Morand C, *et al.* Polyphenols - food sources and bioavailability. *Am J Clin Nutr* 2004; 79:727–747.
A comprehensive review describing the limits of our knowledge on the main classes of polyphenols and key issues for future research.
- 24 Moon J, Tsushida T, Nakahara K, Terao J. Identification of quercetin 3-O-beta-D-glucuronide as an antioxidative metabolite in rat plasma after oral administration of quercetin. *Free Radic Biol Med* 2001; 30:1274–1285.
- 25 Turner R, Baron T, Wolfram S, *et al.* Effect of circulating forms of soy isoflavones on the oxidation of low density lipoprotein. *Free Radic Res* 2004; 38:209–216.
One of the few studies investigating the biological properties of physiological metabolites of polyphenols.
- 26 Stocker R, O'Halloran RA. Dealcoholized red wine decreases atherosclerosis in apolipoprotein E gene-deficient mice independently of inhibition of lipid peroxidation in the artery wall. *Am J Clin Nutr* 2004; 79:123–130.
The authors suggest that dealcoholized red wine contains polyphenolic compounds capable of synergizing with vitamin E, and its long-term moderate consumption can decrease atherosclerosis in apolipoprotein E^{-/-} mice.
- 27 O'Reilly JD, Mallet AI, McAnlis GT, *et al.* Consumption of flavonoids in onions and black tea: lack of effect on F₂-isoprostanes and autoantibodies to oxidized LDL in healthy humans. *Am J Clin Nutr* 2001; 73:1040–1425.
- 28 Hodgson JM, Croft KD, Mori TA, *et al.* Regular ingestion of tea does not inhibit in vivo lipid peroxidation in humans. *J Nutr* 2002; 132:55–58.
- 29 Davies MJ, Judd JT, Baer DJ, *et al.* Black tea consumption reduces total and LDL cholesterol in mildly hypercholesterolemic adults. *J Nutr* 2003; 133:3298S–3302S.
- 30 Cartron E, Fouret G, Carbonneau MA, *et al.* Red-wine beneficial long-term effect on lipids but not on antioxidant characteristics in plasma in a study comparing three types of wine—description of two O-methylated derivatives of gallic acid in humans. *Free Radic Res* 2003; 37:1021–1035.
- 31 Abu-Amsa Caccetta R, Burke V, Mori TA, *et al.* Red wine polyphenols, in the absence of alcohol, reduce lipid peroxidative stress in smoking subjects. *Free Radic Biol Med* 2001; 30:636–642.
- 32 Masella R, Giovannini C, Vari R, *et al.* Effects of dietary virgin olive oil phenols on low density lipoprotein oxidation in hyperlipidemic patients. *Lipids* 2001; 36:1195–1202.
- 33 Marrugat J, Covas MI, Fito M, *et al.* Effects of differing phenolic content in dietary olive oils on lipids and LDL oxidation—a randomized controlled trial. *Eur J Nutr* 2004; 43:140–147.
- 34 Moschandreas J, Vissers MN, Wiseman S, *et al.* Extra virgin olive oil phenols and markers of oxidation in Greek smokers: a randomized cross-over study. *Eur J Clin Nutr* 2002; 56:1024–1029.
- 35 Wan Y, Vinson JA, Etherton TD, *et al.* Effects of cocoa powder and dark chocolate on LDL oxidative susceptibility and prostaglandin concentrations in humans. *Am J Clin Nutr* 2001; 74:596–602.
- 36 Murphy KJ, Chronopoulos AK, Singh I, *et al.* Dietary flavanols and procyanidin oligomers from cocoa (*Theobroma cacao*) inhibit platelet function. *Am J Clin Nutr* 2003; 77:1466–1473.
- 37 Marniemi J, Hakala P, Maki J, Ahotupa M. Partial resistance of low density lipoprotein to oxidation in vivo after increased intake of berries. *Nutr Metab Cardiovasc Dis* 2000; 10:331–337.
- 38 Ventura P, Bini A, Panini R, *et al.* Red wine consumption prevents vascular oxidative stress induced by a high-fat meal in healthy volunteers. *Int J Vitam Nutr Res* 2004; 74:137–143.
- 39 Natella F, Ghiselli A, Guidi A, *et al.* Red wine mitigates the postprandial increase of LDL susceptibility to oxidation. *Free Radic Biol Med* 2001; 30:1036–1044.
- 40 Natella F, Belevi F, Gentili V, *et al.* Grape seed proanthocyanidins prevent plasma postprandial oxidative stress in humans. *J Agric Food Chem* 2002; 50:7720–7725.
- 41 Gimeno E, Fito M, Lamuela-Raventos RM, *et al.* Effect of ingestion of virgin olive oil on human low-density lipoprotein composition. *Eur J Clin Nutr* 2002; 56:114–120.
- 42 Vissers MN, Zock PL, Leenen R, *et al.* Effect of consumption of phenols from olives and extra virgin olive oil on LDL oxidizability in healthy humans. *Free Radic Res* 2001; 35:619–629.
- 43 Weggemans RM, Trautwein EA. Relation between soy-associated isoflavones and LDL and HDL cholesterol concentrations in humans: a meta-analysis. *Eur J Clin Nutr* 2003; 57:940–946.
The selection of the studies used in this meta-analysis and the methods are described very carefully. This analysis provides reliable conclusions on the impact of isoflavones on lipemia.
- 44 Wagner JD, Schwenke DC, Greaves KA, *et al.* Soy protein with isoflavones, but not an isoflavone-rich supplement, improves arterial low-density lipoprotein metabolism and atherogenesis. *Arterioscler Thromb Vasc Biol* 2003; 23:2241–2246.
- 45 Desroches S, Mauger JF, Ausman LM, *et al.* Soy protein favorably affects LDL size independently of isoflavones in hypercholesterolemic men and women. *J Nutr* 2004; 134:574–579.
- 46 Lichtenstein AH, Jalbert SM, Adlercreutz H, *et al.* Lipoprotein response to diets high in soy or animal protein with and without isoflavones in moderately hypercholesterolemic subjects. *Arterioscler Thromb Vasc Biol* 2002; 22:1852–1858.
- 47 Sanders TA, Dean TS, Grainger D, *et al.* Moderate intakes of intact soy protein rich in isoflavones compared with ethanol-extracted soy protein increase HDL but do not influence transforming growth factor beta(1) concentrations and hemostatic risk factors for coronary heart disease in healthy subjects. *Am J Clin Nutr* 2002; 76:373–377.
- 48 Wangen KE, Duncan AM, Xu X, Kurzer MS. Soy isoflavones improve plasma lipids in normocholesterolemic and mildly hypercholesterolemic postmenopausal women. *Am J Clin Nutr* 2001; 73:225–231.
- 49 Blakesmith SJ, Lyons-Wall PM, George C, *et al.* Effects of supplementation with purified red clover (*Trifolium pratense*) isoflavones on plasma lipids and insulin resistance in healthy premenopausal women. *Br J Nutr* 2003; 89:467–474.
- 50 Nestel P, Cehun M, Chronopoulos A, *et al.* A biochanin-enriched isoflavone from red clover lowers LDL cholesterol in men. *Eur J Clin Nutr* 2004; 58:403–408.
- 51 Hashimoto M, Kim S, Eto M, *et al.* Effect of acute intake of red wine on flow-mediated vasodilatation of the brachial artery. *Am J Cardiol* 2001; 88:1457–1460.
- 52 Papamichael C, Karatzis E, Karatzis K, *et al.* Red wine's antioxidants counteract acute endothelial dysfunction caused by cigarette smoking in healthy non-smokers. *Am Heart J* 2004; 147:E5.
- 53 Whelan AP, Sutherland WH, McCormick MP, *et al.* Effects of white and red wine on endothelial function in subjects with coronary artery disease. *Intern Med J* 2004; 34:224–228.
- 54 Chou EJ, Keevil JG, Aeschlimann S, *et al.* Effect of ingestion of purple grape juice on endothelial function in patients with coronary heart disease. *Am J Cardiol* 2001; 88:553–555.
- 55 Vita JA. Tea consumption and cardiovascular disease: effects on endothelial function. *J Nutr* 2003; 133:3293S–3297S.
- 56 Duffy SJ, Keane JF Jr, Holbrook M, *et al.* Short- and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease. *Circulation* 2001; 104:151–156.
- 57 Squadrito F, Altavilla D, Morabito N, *et al.* The effect of the phytoestrogen genistein on plasma nitric oxide concentrations, endothelin-1 levels and endothelium dependent vasodilation in postmenopausal women. *Atherosclerosis* 2002; 163:339–347.
- 58 Cuevas AM, Iribarra VL, Castillo OA, *et al.* Isolated soy protein improves endothelial function in postmenopausal hypercholesterolemic women. *Eur J Clin Nutr* 2003; 57:889–894.
- 59 Engler MB, Engler MM, Chen CY, *et al.* Flavonoid-rich dark chocolate improves endothelial function and increases plasma epicatechin concentrations in healthy adults. *J Am Coll Nutr* 2004; 23:197–204.
- 60 Hodgson JM, Puddey IB, Burke V, *et al.* Regular ingestion of black tea improves brachial artery vasodilator function. *Clin Sci Lond* 2002; 102:195–201.
- 61 Hirata K, Shimada K, Watanabe H, *et al.* Black tea increases coronary flow velocity reserve in healthy male subjects. *Am J Cardiol* 2004; 93:1384–1388.
- 62 Squadrito F, Altavilla D, Crisafulli A, *et al.* Effect of genistein on endothelial function in postmenopausal women: a randomized, double-blind, controlled study. *Am J Med* 2003; 114:470–476.

8 Nutrition and metabolism

- 63** Steinberg FM, Guthrie NL, Villablanca AC, *et al.* Soy protein with isoflavones has favorable effects on endothelial function that are independent of lipid and antioxidant effects in healthy postmenopausal women. *Am J Clin Nutr* 2003; 78:123–130.
- 64** Badia E, Sacanella E, Fernandez-Sola J, *et al.* Decreased tumor necrosis factor- α -induced adhesion of human monocytes to endothelial cells after moderate alcohol consumption. *Am J Clin Nutr* 2004; 80:225–230.
- Tumor necrosis factor- α -induced adhesion of monocytes to endothelial cells was virtually abolished after red-wine consumption in human volunteers but was only partially reduced after gin consumption.
- 65** de Maat MP, Pijl H, Kluit C, Princen HM. Consumption of black and green tea had no effect on inflammation, haemostasis and endothelial markers in smoking healthy individuals. *Eur J Clin Nutr* 2000; 54:757–763.
- 66** Mathur S, Devaraj S, Grundy SM, Jialal I. Cocoa products decrease low density lipoprotein oxidative susceptibility but do not affect biomarkers of inflammation in humans. *J Nutr* 2002; 132:3663–3667.
- 67** Hodgson JM, Puddey IB, Mori TA, *et al.* Effects of regular ingestion of black tea on haemostasis and cell adhesion molecules in humans. *Eur J Clin Nutr* 2001; 55:881–886.
- 68** Teede HJ, McGrath BP, DeSilva L, *et al.* Isoflavones reduce arterial stiffness: a placebo-controlled study in men and postmenopausal women. *Arterioscler Thromb Vasc Biol* 2003; 23:1066–1071.
- 69** Hodgson JM, Puddey IB, Burke V, *et al.* Acute effects of ingestion of black tea on postprandial platelet aggregation in human subjects. *Br J Nutr* 2002; 87:141–145.
- 70** Mezzano D, Leighton F, Martinez C, *et al.* Complementary effects of Mediterranean diet and moderate red wine intake on haemostatic cardiovascular risk factors. *Eur J Clin Nutr* 2001; 55:444–451.
- 71** Mezzano D, Leighton F, Strobel P, *et al.* Mediterranean diet, but not red wine, is associated with beneficial changes in primary haemostasis. *Eur J Clin Nutr* 2003; 57:439–446.
- 72** Rein D, Paglieroni TG, Wun T, *et al.* Cocoa inhibits platelet activation and function. *Am J Clin Nutr* 2000; 72:30–35.
- 73** Hollman PCH, van Trijp JMP, Buysman MNCP, *et al.* Relative bioavailability of the antioxidant flavonoid quercetin from various foods in man. *FEBS Lett* 1997; 418:152–156.
- 74** Meyer BJ, Larkin TA, Owen AJ, *et al.* Limited lipid-lowering effects of regular consumption of whole soybean foods. *Ann Nutr Metab* 2004; 48:67–78.
- Innovative strategy of stratification of the volunteers according to their ability to perform a particular metabolism.
- 75** Zhang Y, Song TT, Cunnick JE, *et al.* Daidzein and genistein glucuronides in vitro are weakly estrogenic and activate human natural killer cells at nutritionally relevant concentrations. *J Nutr* 1999; 129:399–405.
- 76** Cassidy A, De Pascual Teresa S, Rimbach G. Molecular mechanisms by which dietary isoflavones potentially prevent atherosclerosis. *Expert Rev Mol Med* 2003; 5:1–15.
- An excellent and comprehensive review of the various mechanisms whereby isoflavones may prevent atherosclerosis. Bioavailability data are taken into account.

MOL

Manuscript No.

16111

**Current Opinion in Lipidology
Typeset by Thomson Digital for
Lippincott Williams & Wilkins**

Dear Author,

During the preparation of your manuscript for typesetting, some queries have arisen. These are listed below. Please check your typeset proof carefully and mark any corrections in the margin as neatly as possible or compile them as a separate list. This form should then be returned with your marked proof/list of corrections to the Production Editor.

QUERIES: to be answered by AUTHOR/EDITOR

AUTHOR: The following queries have arisen during the editing of your manuscript. Please answer the queries by marking the requisite corrections at the appropriate positions in the text.

QUERY NO.	QUERY DETAILS	
	No Queries	